

# Fitness training

- **Definition:** It consists of planned and structured activities with appropriate intensity, involving continuous rhythmic movements and various muscular groups, aimed at maintaining or increasing components related to physical fitness. It is based upon physiological principles, which increase the ability to transport and use oxygen during physical activity.
- Advantages: Easy administration and relatively low cost. Possibility of being delivered to groups.
- Clearly established evidence for stroke patients with mild to moderate impairments at both the acute and chronic stages in outcomes, such as maximal oxygen consumption, gait capacity and speed.
- Limited scientific evidence to improve quality of life.

## Constraint-induced movement therapy

- **Definition:** It is a behavioural approach, which focuses on improving and increasing the use of the paretic limb in people with asymmetrical use of the limbs. Comprises three pillars: (1) restriction of the non-paretic limb; (2) intensive task-oriented training (task practice and shaping); and (3) a transfer package.
- Advantages: The activities to be trained may be those that are part of the patients' daily lives and be administered within their life context.
- Clearly established evidence for stroke patients at the acute, sub-acute, and chronic stages in outcomes related to the quantity and quality of the use of their paretic upper limb in daily life, dexterity, and motor recovery.
- Limited scientific evidence for the lower limbs, in outcomes related to strength and gait.

# Biofeedback

- **Definition:** It is delivered using equipment to provide patients with information regarding physiological processes during the performance of a given movement, activity, or task, in order to improve performance and learning.
- Advantages: The physiological information is continuously and simultaneously delivered in a objective manner and is specific to the trained movement, activity, or task.
- Clearly established evidence for stroke patients at the acute, sub-acute, and chronic stages to improve activity performances more related to the lower limbs (stand-up, gait, and sit-to-stand).
- Limited scientific evidence for sensorimotor impairments, activities related to the upper limbs, and spasticity that interferes with activity or personal care.

# Electrical stimulation

- **Definition:** It refers to the stimulation of an intact lower motor neuron by an electrical current generated by an apparatus. It can be used to activate paralyzed or paretic muscles, aiming at generating muscular contractions or modulating neural activity, to regain voluntary contractions or to prevent abnormal muscular reactions.
- **Advantages:** In precise sequences and magnitudes, electrical stimulation can be used to directly accomplish functional tasks.
- **Clearly established evidence** for patients at the sub-acute and chronic stages to improve muscular strength and activity performance.
- **Limited scientific evidence** to prevent or treat swelling of the extremities and to deal with spasticity, that interferes with activity or personal care.



# Progressive resistance training

- **Definition:** Refers to the use of progressive overload applied to a specific muscular group, to stimulate further adaptation toward specific training goals.
- **Advantages:** Low cost, easy to administer, and may be delivered in groups. It can be used for muscles of different segments (upper limbs, lower limbs, and trunk) and also for the respiratory muscles.
- **Clearly established evidence** for for weak patients at the acute, sub-acute, and chronic stages, without any adverse effects, even on spasticity, to improve strength, gait performance, quality of life, and oxygen consumption (peak  $\text{VO}_2$ ).
- **Limited scientific evidence** to improve mobility, sit-to-stand, stairclimb, the performance of activities related to the upper limbs, and functional performance in general.



## Mirror therapy

- **Definition:** It focuses on moving the unimpaired limb, while the patient watches its mirror reflection superimposed over the impaired limb that is unseen, thus creating a visual illusion of the movement capability of the impaired limb.
- Advantages: Low cost and is easily performed. The patients may perform the intervention by themselves.
- Clearly established evidence for weak patients at the acute, sub-acute, and chronic stages to improve the following outcomes related to the upper limbs: Pain, motor function, and the performance of functional activities.
- Limited scientific evidence to improve range of motion, visual-spatial neglect, and the performance of activities of daily living.

## Bobath concept

- **Definition:** It is defined by the International Bobath Instructors Training Association (IBITA), as a problem solving approach for the assessment and treatment of individuals with disturbances of function, movement, and postural control due to a lesion of the central nervous system. The aims are to identify and analyze problems within functional activities and participation in everyday life, as well as analyse movement components and underlying impairments. It is based upon afferent information, named facilitation, to enable successful movement and task performance.
- **Advantages:** It is a global strategy with a holistic approach, but it requires expert training to be delivered.
- **Evidence:** There is no evidence that the Bobath Concept is superior to other approaches.

# Transcranial magnetic stimulation

- **Definition:** It is an intervention aiming at altering the excitability of the motor cortex. Improvements in motor performance after stroke have been found after inhibiting the unaffected hemisphere, as well as increasing the excitability of the affected hemisphere.
- Advantages: It can be implemented for stroke patients with severe motor impairments.
- Limited scientific evidence to improve motor recovery at the acute and chronic stages. At the moment, no adverse effects have been reported.



## Task-specific training

- **Definition:** It consists of repetitive training of movements directly related to functional activities.
- **Advantages:** It can be performed as circuit training and delivered to groups. The intensity of training can be adjusted and progressed to fit the individuals' needs.
- **Clearly established evidence** for stroke patients at the sub-acute and chronic stages to improve balance, sit-to-stand, reaching, manipulation, and walking performance.

A person with short grey hair is seen from the back, sitting at a desk and looking at a computer monitor. The monitor displays a virtual environment with a bright, outdoor scene. The person is wearing a light blue long-sleeved shirt. The background is a plain, light-colored wall.

## Virtual reality therapy

- **Definition:** It refers to use of interactive simulations generated from images, such as computer processors, to provide the users the opportunity to interact with environments that simulate real objects and events. Through a virtual environment rich in detail, virtual reality simulates functional tasks that are intensively practiced.
- **Advantages:** Virtual reality programs simulate real life functional activities in an interesting and challenging manner and may encourage the practice of a higher number of repetitions. Moreover, the difficulty of the tasks can be graded and the physiotherapists can simulate tasks that could not be trained within clinical settings, such as crossing a street.
- **Limited scientific evidence** of the benefits of virtual reality compared with the same doses of conventional rehabilitation strategies for measures of upper limb function and daily life activities. There is limited evidence on the effectiveness of virtual reality in measures of grip strength and gait speed.

# Mental practice

- **Definition:** It is a cognitive strategy, in which a brain area related to specific motor action is triggered repeatedly, through the activation of the imagination, with the goal of improving the patients' performance.
- Advantages: For many individuals with nervous system damage, the execution of certain movements is very difficult and, sometimes, even impossible, which hampers their active participation in the rehabilitation process. In this sense, mental practice allows the realization of all movements.
- Limited scientific evidence of the benefits of mental practice in addition to other rehabilitation strategies for measures of upper limb function. No adverse effects with stroke patients at the acute, sub-acute and chronic stages were observed.